arXiv:1511.06661v1 [math.CO] 20 Nov 2015

F-INDEX OF SOME GRAPH OPERATIONS

NILANJAN DE, SK. MD. ABU NAYEEM, AND ANITA PAL

ABSTRACT. The F-index of a graph is defined as the sum of cubes of the vertex degrees of the graph. This was introduced in 1972, in the same paper where the first and second Zagreb indices were introduced to study the structure-dependency of total π-electron energy. But this topological index was not further studied till then. Very recently, Furtula and Gutman [B. Furtula, I. Gutman, A forgotten topological index, *J. Math. Chem.*, **53(4)**(2015) 1184–1190.] reinvestigated the index and named it "forgotten topological index" or "F-index". In that paper, they present some basic properties of this index and showed that this index can enhance the physico-chemical applicability of Zagreb index. Here, we study the behavior of this index under several graph operations and apply our results to find the F-index of different chemically interesting molecular graphs and nano-structures.

Keywords: Topological index; vertex degree; first and second Zagreb indices; F index; graph operations.

Mathematics Subject Classification: Primary: 05C35; Secondary: 05C07, 05C40

1. INTRODUCTION

Suppose G is a simple connected graph and $V(G)$ and $E(G)$ denote the vertex set and edge set of G, respectively. For any vertex $v \in V(G)$, let $d_G(v)$ denote its degree, that is the number of neighbors of v and $N(v)$ denote the set of vertices which are the neighbors of the vertex v, so that $|N(v)| = d_G(v)$. In chemistry, biochemistry and nanotechnology different topological indices are found to be useful in isomer discrimination, structure-property relationship, structure-activity relationship and pharmaceutical drug design. The first and second Zagreb indices of a graph G, denoted by $M_1(G)$ and $M_2(G)$, are among the oldest, most popular and extremely studied vertex-degree based topological indices and are respectively defined as

$$M_1(G) = \sum_{v \in V(G)} d_G(v)^2 = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]$$

and

$$M_2(G) = \sum_{uv \in E(G)} d_G(u) d_G(v).$$

These indices were introduced in a paper in 1972 [1] to study the structure-dependency of the total π-electron energy (ε). It was found that the ε depends on $M_1(G)$ and thus provides a measure of carbon skeleton of the underlying molecules. In the same paper, another topological index, defined as sum of cubes of degrees of the vertices of the graph was also shown to influence ε. However this index was not further studied till then, except in a recent article by Furtula and Gutman [2] where they reinvestigated this index and studied some basic properties of this index. They showed that the predictive ability of this index is almost similar to that of first Zagreb index and for the entropy and acentric factor, both of them yield correlation coefficients greater than 0.95. They named this index as "forgotten topological index" or "F-index". Throughout the present paper we name this index as F-index and denote it by $F(G)$, so that

$$F(G) = \sum_{v \in V(G)} d_G(v)^3 = \sum_{uv \in E(G)} [d_G(u)^2 + d_G(v)^2].$$

As we know that some chemically interesting graphs can be obtained by different graph operations on some general or particular graphs, it is important to study such graph operations in order to understand how it is related to the corresponding topological indices of the original graphs. In [3], Khalifeh et al. derived some exact formulae for computing first and second Zagreb indices under some graph operations. In [4], Das et al. derived some upper bounds for multiplicative Zagreb indices for different graph operations. In [5], Azari presented some lower bounds for Narumi-Katayama index under several graph operations. In [10], the present authors computed some bounds and exact formulae of the connective eccentric index under different graph operations. There are several other results regarding various topological indices under different graph operations are available in the literature. In [6], Azari and Iranmanesh presented explicit formulas for computing the eccentric-distance sum of different graph operations. Interested readers are referred to [11, 12, 13, 14, 15, 8, 7, 9] in this regard.

In this paper, we present some exact expressions for the F-index of different graph operations such as union, join, Cartesian product, composition, tensor product, strong product, corona product, generalized hierarchical product, disjunction, symmetric difference, splice and link of graphs. Also we apply our results to compute the F-index for some important classes of molecular graphs and nano-structures.

2. Main Results and Discussions

In this section, we study F-index of various graph operations like union, join, Cartesian product, composition, tensor product, strong product, corona product, generalized hierarchical product, disjunction, symmetric difference, link and splice of graphs. These operations are binary and all the graphs are connected, finite and simple. In the following, if not indicated otherwise, we use the notation $V(G_i)$ for the vertex set, $E(G_i)$ for the edge set, n_i for the number of vertices and m_i for the number of edges of the graph G_i, $i \in \{1,2,\ldots,k\}$, respectively. Throughout the paper, we use the familiar notations P_n, C_n and K_n to denote a path graph, cycle graph and complete graph with n number of vertices, respectively.

2.1. **Union.** Let G_1, G_2,...,G_k be k graphs with disjoint vertex sets. Then their union $G_1 \cup G_2 \cup ... \cup G_k$ is the graph with vertex set $V(G_1) \cup V(G_2) \cup ... \cup V(G_k)$ and the edge set $E(G_1) \cup E(G_2) \cup ... \cup E(G_k)$. The degree of a vertex v of $G_1 \cup G_2 \cup ... \cup G_k$ is equal to the degree of the vertex v in the component G_i, $i = 1,2,...,k$, that contains it. In the following theorem we obtain the F-index of the union of k number of graphs.

Theorem 1. *The F-index of $G_1 \cup G_2 \cup ... \cup G_k$ is given by*

$$F(G_1 \cup G_2 \cup ... \cup G_k) = F(G_1) + F(G_2) + ... + F(G_k).$$

Proof. By definition of F-index we have

$$\begin{aligned} F(G_1 \cup G_2 \cup ... \cup G_k) &= \sum_{v \in V(G_1 \cup G_2 \cup ... \cup G_k)} (d_{G_1 \cup G_2 \cup ... \cup G_k}(v))^3 \\ &= \sum_{v \in V(G_1)} d_{G_1}(v)^3 + \sum_{v \in V(G_2)} d_{G_2}(v)^3 + ... + \sum_{v \in V(G_k)} d_{G_k}(v)^3 \\ &= F(G_1) + F(G_2) + ... + F(G_k), \end{aligned}$$

which completes the proof. □

2.2. **Join.** The join $G_1 + G_2$ of two graphs G_1 and G_2 is the union $G_1 \cup G_2$ together with all the edges joining $V(G_1)$ and $V(G_2)$. The degree of a vertex v of $G_1 + G_2$ is

$$d_{G_1+G_2}(v) = \begin{cases} d_{G_1}(v) + n_2, v \in V(G_1) \\ d_{G_2}(v) + n_1, v \in V(G_2). \end{cases}$$

In general, for k graphs G_1, G_2,...,G_k, the degree of a vertex v in $G_1 + G_2 + ... + G_k$ is given by $d_{G_1+G_2+...+G_k}(v) = d_{G_i}(v) + n - n_i$, where v is originally a vertex of the graph G_i and $n = n_1 + n_2 + ... + n_k$.

In the following theorem we compute the F-index of the join of k number of graphs.

Theorem 2. *The F-index of $G_1 + G_2 + ... + G_k$ is given by*

$$F(G_1 + G_2 + ... + G_k) = \sum_{i=1}^{k} F(G_i) + 3\sum_{i=1}^{k} \bar{n}_i M_1(G_i) + 6\sum_{i=1}^{k} \bar{n}_i^2 m_i + \sum_{i=1}^{k} n_i \bar{n}_i^3,$$

where $\bar{n}_i = n - n_i, i = 1,2,...,k$ and $n = n_1 + n_2 + ... + n_k$.

Proof. We have

$$\begin{aligned} F(G_1 + G_2 + ... + G_k) &= \sum_{i=1}^{k} \sum_{v \in V(G_i)} (d_{G_i}(v) + \bar{n}_i)^3 \\ &= \sum_{i=1}^{k} \sum_{v \in V(G_i)} (d_{G_i}(v)^3 + 3\bar{n}_i d_{G_i}(v)^2 + 3\bar{n}_i^2 d_{G_i}(v) + \bar{n}_i^3) \\ &= \sum_{i=1}^{k} \sum_{v \in V(G_i)} d_{G_i}(v)^3 + 3\sum_{i=1}^{k} \bar{n}_i \sum_{v \in V(G_i)} d_{G_i}(v)^2 \\ &\quad + 3\sum_{i=1}^{k} \bar{n}_i^2 \sum_{v \in V(G_i)} d_{G_i}(v) + \sum_{i=1}^{k} n_i \bar{n}_i^3, \end{aligned}$$

which completes the proof. □

Let $G_1 = G_2 = ... = G_p = G$ and pG denote the join of p copies of G. Then the following corollaries follow as direct consequence of the previous theorem.

Corollary 1. *Let, n and m be the number of vertices and edges of G, respectively. Then*

$$F(pG) = pF(G) + 3np(p-1)M_1(G) + 6n^2 mp(p-1)^2 + n^4 p(p-1)^3.$$

Corollary 2. *The F-index of $G_1 + G_2$ is given by*

$$F(G_1 + G_2) = F(G_1) + F(G_2) + 3n_2 M_1(G_1) + 3n_1 M_1(G_2) + 6{n_2}^2 m_1 + 6{n_1}^2 m_2 + n_1 {n_2}^3 + n_2 {n_1}^3.$$

The suspension of a graph G is defined as $K_1 + G$. So from the Corollary 2 the following result follows.

Corollary 3. *Let, n and m be the number of vertices and edges of G, respectively. Then the F-index of suspension of G is given by*

$$F(K_1 + G) = F(G) + 3M_1(G) + n^3 + 6m + n.$$

Example 1. *The complete n-partite graph $K_{m_1,m_2,...,m_n}$ (Fig.1) on $m_1 + m_2 + ... + m_n$ vertices can be considered as $\bar{K}_{m_1} + \bar{K}_{m_2} + ... + \bar{K}_{m_n}$. Then the F-index of $K_{m_1,m_2,...,m_n}$ is given by*

$$F(K_{m_1,m_2,...,m_n}) = \sum_{i=1}^{n} m_i \bar{m}_i^3,$$

where $\bar{m}_i = (m_1 + m_2 + ... + m_n) - m_i, i = 1,2,...,n.$



FIGURE 1. The complete n-partite graph.

Example 2. *The wheel graph W_n on $(n + 1)$ vertices is the suspension of C_n and the fan graph F_n on $(n + 1)$ vertices is the suspension of P_n. So their F-indices are given by*

(i) $F(W_n) = n^3 + 27n$,

(ii) $F(F_n) = n^3 + 27n - 38$.

Example 3. *The dutch windmill graph or flower graph is the suspension of m copies of K_2, denoted by mK_2. So its F-index is given by $F(K_1 + mK_2) = 8m^3 + 16m$.*

Example 4. *The cone graph $C_{m,n}$ is defined as $C_m + \bar{K}_n$. So its F-index is calculated as $F(C_{m,n}) = mn^3 + m^3n + 6mn^2 + 12mn + 8m$.*

2.3. **Cartesian Product.** The Cartesian product of G_1 and G_2, denoted by $G_1 \otimes G_2$, is the graph with vertex set $V(G_1) \times V(G_2)$ and any two vertices (u_p, v_r) and (u_q, v_s) are adjacent if and only if $[u_p = u_q$ and $v_r v_s \in E(G_2)]$ or $[v_r = v_s$ and $u_p u_q \in E(G_1)]$. In the following theorem first we find the first Zagreb index of the Cartesian product of k number of graphs.

Theorem 3. *Let n be the total number of vertices in $\bigotimes_{i=1}^{k} G_i$, then the first Zagreb index of $G_1 \otimes G_2 \otimes ... \otimes G_k$ is given by*

$$M_1\left(\bigotimes_{i=1}^{k} G_i\right) = n\sum_{i=1}^{k}\frac{M_1(G_i)}{n_i} + 4n\sum_{\substack{i,j=1\\ i\neq j}}^{k}\frac{m_i}{n_i}\cdot\frac{m_j}{n_j}.$$

Proof. For the proof of this theorem we refer to Theorem 1 of [3]. □

In the following theorem we obtain the F-index of the Cartesian product of k number of graphs.

Theorem 4. *The F-index of $G_1 \otimes G_2 \otimes ... \otimes G_k$ is given by*

$$F(G_1 \otimes G_2 \otimes ... \otimes G_k) = n\sum_{i=1}^{k}\frac{F(G_i)}{n_i} + 6n\sum_{\substack{i,j=1\\ i\neq j}}^{k}\frac{M_1(G_i)}{n_i}\cdot\frac{m_j}{n_j} + 8n\sum_{\substack{p,q,r=1\\ p\neq q\neq r}}^{k}\frac{m_p}{n_p}\cdot\frac{m_q}{n_q}\cdot\frac{m_r}{n_r}.$$

Proof. First we prove the result for $k = 2$.

We have, $d_{G_1\otimes G_2}(a,b) = d_{G_1}(a) + d_{G_2}(b)$. So, from definition of F-index we have

$$\begin{aligned}
F(G_1 \otimes G_2) &= \sum_{a\in V(G_1)}\sum_{b\in V(G_2)}\left(d_{G_1}(a) + d_{G_2}(b)\right)^3 \\
&= \sum_{b\in V(G_2)}\sum_{a\in V(G_1)} d_{G_1}(a)^3 + \sum_{a\in V(G_1)}\sum_{b\in V(G_2)} d_{G_2}(b)^3 \\
&\quad +3\sum_{a\in V(G_1)} d_{G_1}(a)^2 \sum_{b\in V(G_2)} d_{G_2}(b) + 3\sum_{a\in V(G_1)} d_{G_1}(a)\sum_{b\in V(G_2)} d_{G_2}(b)^2 \\
&= n_2F(G_1) + n_1F(G_2) + 6m_2M_1(G_1) + 6m_1M_1(G_2).
\end{aligned}$$

Let n' be the total number of vertices in $\bigotimes_{i=1}^{k-1} G_i$. Then by an inductive argument, using Theorem 3, we have

$$\begin{aligned}
F\left(\bigotimes_{i=1}^{k} G_i\right) &= F\left(\bigotimes_{i=1}^{k-1} G_i \bigotimes G_k\right) \\
&= n_k F\left(\bigotimes_{i=1}^{k-1} G_i\right) + n' F(G_k) + 6m_k M_1\left(\bigotimes_{i=1}^{k-1} G_i\right) + 6n' \sum_{i=1}^{k-1} \frac{m_i}{n_i} M_1(G_k) \\
&= n_k \left[n' \sum_{i=1}^{k-1} \frac{F(G_i)}{n_i} + 6n' \sum_{\substack{i,j=1 \\ i\neq j}}^{k-1} \frac{M_1(G_i)}{n_i} \cdot \frac{m_j}{n_j} + 8n' \sum_{\substack{p,q,r=1 \\ p\neq q\neq r}}^{k-1} \frac{m_p}{n_p} \cdot \frac{m_q}{n_q} \cdot \frac{m_r}{n_r} \right] \\
&\quad + n' F(G_k) + 6m_k \left[n' \sum_{i=1}^{k-1} \frac{M_1(G_i)}{n_i} + 4n' \sum_{\substack{i,j=1 \\ i\neq j}}^{k-1} \frac{m_i}{n_i} \cdot \frac{m_j}{n_j} \right] + 6n' M_1(G_k) \sum_{i=1}^{k-1} \frac{m_i}{n_i} \\
&= \left[n \sum_{i=1}^{k-1} \frac{F(G_i)}{n_i} + n \frac{F(G_k)}{n_k} \right] + \left[6n \sum_{\substack{i,j=1 \\ i\neq j}}^{k-1} \frac{M_1(G_i)}{n_i} \cdot \frac{m_j}{n_j} + 6n \frac{m_k}{n_k} \sum_{i=1}^{k-1} \frac{M_1(G_i)}{n_i} \right. \\
&\quad \left. + 6n \frac{M_1(G_k)}{n_k} \sum_{i=1}^{k-1} \frac{m_i}{n_i} \right] + \left[8n \sum_{\substack{p,q,r=1 \\ p\neq q\neq r}}^{k-1} \frac{m_p}{n_p} \cdot \frac{m_q}{n_q} \cdot \frac{m_r}{n_r} + 24n \frac{m_k}{n_k} \sum_{\substack{i,j=1 \\ i\neq j}}^{k-1} \frac{m_i}{n_i} \cdot \frac{m_j}{n_j} \right],
\end{aligned}$$

from where the desired result follows. □

Corollary 4. *The F-index of of the torus $C_{n_1} \otimes C_{n_2} \otimes ... \otimes C_{n_k}$ is given by*
$F(C_{n_1} \otimes C_{n_2} \otimes ... \otimes C_{n_k}) = 8k^3 n_1 n_2 \dots n_k$.

Corollary 5. *The F-index of the Hamming graph $K_{n_1} \otimes K_{n_2} \otimes ... \otimes K_{n_k}$ is given by*

$$\begin{aligned}
F(K_{n_1} \otimes K_{n_2} \otimes ... \otimes K_{n_k}) &= n_1 n_2 \dots n_k \Big(\sum_{i=1}^{k} (n_i - 1)^3 + 3 \sum_{\substack{i,j=1 \\ i\neq j}}^{k} (n_i - 1)^2 (n_j - 1) \\
&\quad + \sum_{\substack{p,q,r=1 \\ p\neq q\neq r}}^{k} (n_p - 1)(n_q - 1)(n_r - 1) \Big) \\
&= (n_1 + n_2 + ... + n_k - k)^3 n_1 n_2 ... n_k.
\end{aligned}$$

Example 5. *For k-dimensional hypercube $Q_k = K_2 \otimes K_2 \otimes ... \otimes K_2$ (k times), by our calculation, we have $F(Q_k) = 2^k k^3$.*

Example 6. *The F-index of $K_{n_1} \otimes K_{n_2}$ torus is given by*

$$F(K_{n_1} \otimes K_{n_2}) = n_1 n_2 (n_1 + n_2 - 2)^3$$

Example 7. *Let R and S denote a C_4 nanotube and nanotorus, respectively. Then $R \cong P_n \otimes C_m$ and $S \cong C_n \otimes C_m$, for some integers n and m. Then by our calculation, $F(R) = 64mn - 74m$ and $F(S) = 64mn$. Also if, $T \cong P_n \otimes P_m$, then $F(T) = 64mn - 74m - 74n + 72$.*

2.4. **Composition.** The composition or lexicographic product of two graphs G_1 and G_2 is denoted by $G_1[G_2]$. The vertex set of $G_1[G_2]$ is $V(G_1) \times V(G_2)$ and the degree of a vertex (a,b) of $G_1[G_2]$ is given by $d_{G_1[G_2]}(a,b) = n_2 d_{G_1}(a) + d_{G_2}(b)$ and any two vertices (u_1,u_2) and (v_1,v_2) are adjacent if and only if $u_1v_1 \in E(G_1)$ or $[u_1 = v_1$ and $u_2v_2 \in E(G_2)]$. In the following theorem we compute the F-index of the composition of two graphs.

Theorem 5. *The F-index of $G_1[G_2]$ is given by*

$$F(G_1[G_2]) = {n_2}^4 F(G_1) + n_1 F(G_2) + 6{n_2}^2 m_2 M_1(G_1) + 6n_2 m_1 M_1(G_2).$$

Proof. From definition of F-index, we have

$$\begin{aligned} F(G_1[G_2]) &= \sum_{(a,b)\in V(G_1[G_2])} d_{G_1[G_2]}(a,b)^3 \\ &= \sum_{a\in V(G_1)} \sum_{b\in V(G_2)} [n_2 d_{G_1}(a) + d_{G_2}(b)]^3 \\ &= {n_2}^3 \sum_{b\in V(G_2)} \sum_{a\in V(G_1)} d_{G_1}(a)^3 + \sum_{a\in V(G_1)} \sum_{b\in V(G_2)} d_{G_2}(b)^3 \\ &\quad +3{n_2}^2 \sum_{a\in V(G_1)} d_{G_1}(a)^2 \sum_{b\in V(G_2)} d_{G_2}(b) + 3n_2 \sum_{a\in V(G_1)} d_{G_1}(a) \sum_{b\in V(G_2)} d_{G_2}(b)^2 \\ &= {n_2}^4 F(G_1) + n_1 F(G_2) + 6{n_2}^2 m_2 M_1(G_1) + 6n_2 m_1 M_1(G_2), \end{aligned}$$

which completes proof. □

Example 8. *The fence graph is the composition of P_n and P_2 and the closed fence graph is the composition of C_n and P_2. So from the previous theorem, we have*

(i) $F(P_n[P_2]) = 250n - 392$,

(ii) $F(C_n[P_2]) = 250n$.

2.5. **Tensor Product.** The tensor product or Kronecker product of two graphs G_1 and G_2 is denoted by $G_1 \times G_2$ and any two vertices (u_1, v_1) and (u_2, v_2) are adjacent if and only if $u_1u_2 \in E(G_1)$ and $v_1v_2 \in E(G_2)$. The degree of a vertex (a, b) of $G_1 \times G_2$ is given by $d_{G_1 \times G_2}(a, b) = d_{G_1}(a)d_{G_2}(b)$. In the following theorem, the F-index of the tensor product of two graphs is computed.

Theorem 6. *The F-index of $G_1 \times G_2$ is given by $F(G_1 \times G_2) = F(G_1)F(G_2)$.*

Proof. From definition of F-index, we have

$$F(G_1 \times G_2) = \sum_{(a,b) \in V(G_1 \times G_2)} d_{G_1 \times G_2}(a, b)^3 = \sum_{a \in V(G_1)} \sum_{b \in V(G_2)} [d_{G_1}(a)d_{G_2}(b)]^3 = F(G_1)F(G_2).$$

□

Example 9. *(i)* $F(P_n \times P_m) = (8n - 14)(8m - 14)$,
(ii) $F(C_n \times C_m) = 64nm$,
(iii) $F(K_n \times K_m) = nm(n - 1)^3(m - 1)^3$,
(iv) $F(P_n \times C_m) = 8m(8n - 14)$,
(v) $F(P_n \times K_m) = m(8n - 14)(m - 1)^3$,
(vi) $F(C_n \times K_m) = 8nm(m - 1)^3$.

2.6. **Strong Product.** The strong product of two graphs G_1 and G_2 is denoted by $G_1 \boxtimes G_2$. It has the vertex set $V(G_1) \times V(G_2)$ and any two vertices (u_p, v_r) and (u_q, v_s) are adjacent if and only if $[u_p = u_q$ and $v_rv_s \in E(G_2)]$ or $[v_r = v_s$ and $u_pu_q \in E(G_1)]$ or $[u_pu_q \in E(G_1)$ and $v_rv_s \in E(G_2)]$. Note that if both G_1 and G_2 are connected then $G_1 \boxtimes G_2$ is also connected. The degree of a vertex (a, b) of $G_1 \boxtimes G_2$ is given by

$$d_{G_1 \boxtimes G_2}(a, b) = d_{G_1}(a) + d_{G_2}(b) + d_{G_1}(a)d_{G_2}(b).$$

In the following theorem we compute the F-index of the strong product of two graphs.

Theorem 7. *The F-index of $G_1 \boxtimes G_2$ is given by*

$$F(G_1 \boxtimes G_2) = n_2F(G_1) + n_1F(G_2) + F(G_1)F(G_2) + 6m_2M_1(G_1) + 6m_1M_1(G_2) + 6m_2F(G_1)$$

$$+6m_1F(G_2) + 3F(G_2)M_1(G_1) + 3F(G_1)M_1(G_2) + 6M_1(G_1)M_1(G_2).$$

Proof. From definition of F-index, we have

$$\begin{aligned} F(G_1 \boxtimes G_2) &= \sum_{(v_1,v_2)\in V(G_1\otimes G_2)} d_{G_1\otimes G_2}(v_1,v_2)^3 \\ &= \sum_{v_1\in V(G_1)}\sum_{v_2\in V(G_2)} [d_{G_1}(v_1)+d_{G_2}(v_2)+d_{G_1}(v_1)d_{G_2}(v_2)]^3 \\ &= \sum_{v_1\in V(G_1)}\sum_{v_2\in V(G_2)} [d_{G_1}(v_1)^3+d_{G_2}(v_2)^3+d_{G_1}(v_1)^3d_{G_2}(v_2)^3+3d_{G_1}(v_1)^2d_{G_2}(v_2) \\ &\quad +3d_{G_1}(v_1)d_{G_2}(v_2)^2+3d_{G_1}(v_1)^3d_{G_2}(v_2)+3d_{G_1}(v_1)^3d_{G_2}(v_2)^2 \\ &\quad +3d_{G_2}(v_2)^3d_{G_1}(v_1)+3d_{G_1}(v_1)^2d_{G_2}(v_2)^3+6d_{G_1}(v_1)^2d_{G_2}(v_2)^2]. \end{aligned}$$

On simplification, we obtain the desired result. □

2.7. **Corona Product.** The corona product $G_1 \odot G_2$ of two graphs G_1 and G_2 is obtained by taking one copy of G_1 and n_1 copies of G_2 and by joining each vertex of the i-th copy of G_2 to the i-th vertex of G_1, where $1 \leq i \leq n_1$. The corona product of G_1 and G_2 has total $(n_1n_2 + n_1)$ number of vertices and $(m_1 + n_1m_2 + n_1n_2)$ number of edges. Clearly, the corona product of two graphs is not commutative. Different topological indices under the corona product of graphs have already been studied by some researchers [21, 23]. It is easy to see that the degree of a vertex v of $G_1 \odot G_2$ is given by

$$d_{G_1\odot G_2}(v) = \begin{cases} d_{G_1}(v)+n_2, v\in V(G_1) \\ d_{G_2}(v)+1, v\in V(G_{2,i}), i=1,2,\ldots,n_1. \end{cases}$$

where, $G_{2,i}$ is the i-th copy of the graph G_2. In the following theorem, the F-index of the corona Product of two graphs is computed.

Theorem 8. *The F-index of $G_1 \odot G_2$ is given by*

$$F(G_1 \odot G_2) = F(G_1) + n_1F(G_2) + 3n_2M_1(G_1) + 3n_1M_1(G_2) + 6{n_2}^2m_1 + 6n_1m_2 + n_1n_2({n_2}^2+1).$$

Proof. From definition of F-index, we have

$$\begin{aligned}
F(G_1 \odot G_2) &= \sum_{v\in V(G_1)} (d_{G_1}(v)+n_2)^3 + n_1 \sum_{v\in V(G_2)} (d_{G_2}(v)+1)^3 \\
&= \sum_{v\in V(G_1)} (d_{G_1}(v)^3 + 3n_2 d_{G_1}(v)^2 + 3{n_2}^2 d_{G_1}(v) + {n_2}^3) \\
&\quad + n_1 \sum_{v\in V(G_2)} (d_{G_2}(v)^3 + 3d_{G_2}(v)^2 + 3d_{G_2}(v) + 1) \\
&= \sum_{v\in V(G_1)} d_{G_1}(v)^3 + 3n_2 \sum_{v\in V(G_1)} d_{G_1}(v)^2 + 3{n_2}^2 \sum_{v\in V(G_1)} d_{G_1}(v) + n_1{n_2}^3 \\
&\quad + n_1 \sum_{v\in V(G_1)} d_{G_2}(v)^3 + 3n_1 \sum_{v\in V(G_1)} d_{G_2}(v)^2 + 3n_1 \sum_{v\in V(G_1)} d_{G_2}(v) + n_1 n_2 \\
&= F(G_1) + 3n_2 M_1(G_1) + 6{n_2}^2 m_1 + n_1{n_2}^3 + n_1 F(G_2) + 3n_1 M_1(G_2) \\
&\quad + 6n_1 m_2 + n_1{n_2}^3 + n_1 n_2.
\end{aligned}$$

□

The t-thorny graph G^t of a given graph G is obtained by joining t-number of thorns (pendent edges) to each vertex of G. A variety of topological indices of thorn graphs have been studied by a number of researchers [22, 24, 25, 26].It is well known that, the t-thorny graph of G is defined as the corona product of G and complement of complete graph with t vertices $\bar{K}_t$. Thus from the previous theorem the following corollary follows.

Corollary 6. *The F-index of the t-thorny graph is given by*

$$F(G^t) = F(G) + 3tM_1(G) + 6mt^2 + nt^3 + nt$$

where, n and m are number of vertices and edges of G, respectively.

Example 10. *The F-index of t-thorny graph of C_n and P_n are given by*

(i) $F({C_n}^t) = nt^3 + 6nt^2 + 13nt + 8n$,

(ii) $F({P_n}^t) = nt^3 + 6nt^2 - 6t^2 + 13nt - 18t + 8n - 14$.

Example 11. *One of the hydrogen suppressed molecular graph is the bottleneck graph (B) of a graph G, which is defined as the corona product of K_2 and G, where G is a given graph. The F-index of bottleneck graph of G is given by $F(B) = 2F(G) + 6M_1(G) + 2n^3 + 6n^2 + 8n + 12m + 2$, where n and m are the number of vertices and edges of G, respectively.*

Next, as an application of corona product of graphs, we find the F index of some particular bridge graphs. Let $G_1, G_2, ..., G_n$ be a set of finite pairwise disjoint graphs. The bridge graph with respect to the vertices $v_1, v_2, ..., v_n$, denoted by $B(G_1, G_2, ..., G_n; v_1, v_2, ..., v_n)$ is the graph obtained by connecting the vertex v_i of G_i and the vertex v_{i+1} of G_{i+1} by an edge for all $i = 1, 2, ..., n-1$. If $G_1 \cong G_2 \cong ... \cong G_n$ and $v_1 = v_2 = ... = v_n = v$, we define $G_n(G, v) = B(G, G, ..., G; v, v, ..., v)$. In particular, let $B_n = G_n(P_3, v)$, where v is the degree 2 vertex of P_3 and $T_{n,k} = G_n(C_k, u)$ be special bridge graphs. Then from definition of corona product of graphs, $B_n \cong P_n \odot \overline{K}_2$ and $T_{n,3} \cong P_n \odot K_2$. Using the previous theorem the F index of these bridge graphs are calculated as follows.

Example 12. *(i)* $F(B_m) = 66m - 74$, *for* $m \geq 2$

(ii) $F(T_{m,3}) = 80m - 74$, *for* $m \geq 2$

(iii) $F(J_{n,m+1}) = n^3m + 6n^2m - 6n^2 + 39nm + 8m - 18n - 14$, *for* $n \geq 3$ *and* $m \geq 2$.

2.8. **Generalized Hierarchical Product.** As an extension of Cartesian product of graphs, Barriere et. al. introduced the generalized hierarchical product of graphs in 2009 [16]. Several results on different topological indices under generalized hierarchical product of graphs are already studied [17, 18, 19, 20]. Let G_1 and G_2 be two connected graphs and $\phi \neq U \subseteq V(G_2)$. Then the hierarchical product of G_1 and G_2 denoted by $G_1 \Pi G_2(U)$, is the graph with vertex set $V(G_1) \times V(G_2)$ and any two vertices (u, v) and (u', v') of $G_1(U)\Pi G_2$ are adjacent if and only if $[u = u' \in V(G_1)$ and $vv' \in E(G_2)]$ or $[v = v' \in U$ and $uu' \in E(G_1)]$.

Clearly, the degree of a vertex (u_1, u_2) of $G_1 \Pi G_2(U)$ is given by

$$d_{G_1 \Pi G_2(U)}(u) = \begin{cases} d_{G_1}(u_1) + d_{G_2}(u_2), u_2 \in U \\ d_{G_2}(u_2), u_2 \in V(G_2) - U. \end{cases}$$

In the following theorem we compute the F-index of the generalized hierarchical product of two graphs.

Theorem 9. *The F index of* $G_1 \Pi G_2(U)$ *is given by*

$$F(G_1 \Pi G_2(U)) = |U| F(G_1) + n_1 F(G_2) + 3M_1(G_1) \sum_{v \in U} d_{G_2}(v) + 6m_1 \sum_{v \in U} d_{G_2}(v)^2.$$

Proof. We have, from definition of F index

$$\begin{aligned} F(G_1\Pi G_2(U)) &= \sum_{(v_1,v_2)\in V(G_1\Pi G_2(U))} d_{G_1\Pi G_2(U)}(v_1,v_2)^3 \\ &= \sum_{v_1\in V(G_1)}\sum_{v_2\in U}[d_{G_1}(v_1)+d_{G_2}(v_2)]^3 + \sum_{v_1\in V(G_1)}\sum_{v_2\in V(G_2)-U} d_{G_2}(v_2)^3 \\ &= \sum_{v_1\in V(G_1)}\sum_{v_2\in U}[d_{G_1}(v_1)^3+d_{G_2}(v_2)^3+3d_{G_1}(v_1)^2d_{G_2}(v_2)+3d_{G_1}(v_1)d_{G_2}(v_2)^2] \\ &\quad + \sum_{v_1\in V(G_1)}\sum_{v_2\in V(G_2)-U} d_{G_2}(v_2)^3 \\ &= |U|F(G_1)+n_1F(G_2)+3M_1(G_1)\sum_{v\in U} d_{G_2}(v)+6m_1\sum_{v\in U} d_{G_2}(v)^2, \end{aligned}$$

which completes the proof. □

From definition of the Cartesian product of graphs, it is clear that the Cartesian product of graphs is a special case of generalized hierarchical product of graphs, that is, if $U = V(G_2)$, then $G_1\Pi G_2(U) \cong G_1 \otimes G_2$. So from the previous theorem we can also obtain the Theorem 3, for $k = 2$.

The cluster product of two graphs G_1 and G_2, denoted by $G_1\{G_2\}$, is obtained by taking one copy of G_1 and n_1 copies of a rooted graph G_2 and by identifying the root of the i-th copy of G_2 with the i-th vertex of G_1, $i = 1,2,...,n_1$. From definition of cluster product of graphs, $|V(G_1\{G_2\})| = n_1n_2$ and $|E(G_1\{G_2\})| = (m_2 + n_1m_2)$. Suppose the root vertex of G_2 is denoted by x. Note that, if $U = \{x\} \subset V(G_1)$ then $G_1\{G_2\} \cong G_1\Pi G_2(U) \cong G_1\Pi G_2(\{x\})$. Then from the previous theorem the following results follow.

Corollary 7. *Let G_1 and G_2 be two connected graphs and x be the root vertex of G_2, then*

$$F(G_1\{G_2\}) = F(G_1) + n_1F(G_2) + 3M_1(G_1)d_{G_2}(x) + 6m_1d_{G_2}(x)^2.$$

Note that, if $U = \{x\}$, x is the root vertex of the graph G_2, then $G_1\Pi G_2(U) = G_1\Pi G_2$, the (standard) hierarchical product of two graphs. It is easy to see that $G_1\Pi G_2 \cong G_1\{G_2\}$.

Example 13. *The square comb lattice $Cq(N)$ with $N = n^2$ vertices can be represented as the cluster product $P_n\{P_n\}$, where the root of P_n is one of its vertices of degree 1. Then from the previous corollary, the F-index of $Cq(N)$ is given by $F(Cq(N)) = 8n^2 + 12n - 38$.*

Example 14. *The sun graph is defined as $Sun_{(m,n)} = C_m\{P_{n+1}\}$, such that P_{n+1} is rooted at a vertex of degree one. Then the F-index of $Sun_{(m,n)}$ is given by $Sun_{(m,n)} = 4m(2n + 5)$.*

2.9. **Disjunction.** The disjunction of two graphs G_1 and G_2, denoted by $G_1 \wedge G_2$, consists of the vertex set $V(G_1) \times V(G_2)$ and two vertices (u_1, v_1) and (u_2, v_2) are adjacent whenever $u_1u_2 \in E(G_1)$ or $v_1v_2 \in E(G_2)$. Clearly, the degree of a vertex (u_1, u_2) of $G_1 \wedge G_2$ is given by

$$d_{G_1 \wedge G_2}(u_1, u_2) = n_1 d_{G_1}(u_1) + n_2 d_{G_2}(u_2) - d_{G_1}(u_1) d_{G_2}(u_2).$$

In the following theorem we obtain the F-index of the disjunction of two graphs.

Theorem 10. *The F-index of $G_1 \wedge G_2$ is given by*

$$\begin{aligned} F(G_1 \wedge G_2) &= {n_2}^4 F(G_1) + {n_1}^4 F(G_2) - F(G_1)F(G_2) + 6n_1{n_2}^2 m_2 M_1(G_1) + 6{n_1}^2 n_2 m_1 M_1(G_2) \\ &\quad + 3n_2 F(G_1) M_1(G_2) + 3n_1 F(G_2) M_1(G_1) - 6{n_2}^2 m_2 F(G_1) - 6{n_1}^2 m_1 F(G_2) \\ &\quad - 6n_1 n_2 M_1(G_1) M_1(G_2). \end{aligned}$$

Proof. We have, from definition of F-index

$$\begin{aligned} F(G_1 \wedge G_2) &= \sum_{(u_1,u_2) \in V(G_1 \wedge G_2)} d_{G_1 \wedge G_2}(u_1, u_2)^3 \\ &= \sum_{u_1 \in V(G_1)} \sum_{u_2 \in V(G_2)} (n_2 d_{G_1}(u_1) + n_1 d_{G_2}(u_2) - d_{G_1}(u_1) d_{G_2}(u_2))^3 \\ &= \sum_{u_1 \in V(G_1)} \sum_{u_2 \in V(G_2)} ({n_2}^3 d_{G_1}(u_1)^3 + {n_1}^3 d_{G_2}(u_2)^3 - d_{G_1}(u_1)^3 d_{G_2}(u_2)^3 \\ &\quad + 3n_1{n_2}^2 d_{G_1}(u_1)^2 d_{G_2}(u_2) + 3{n_1}^2 n_2 d_{G_1}(u_1) d_{G_2}(u_2)^2 - 3{n_2}^2 d_{G_1}(u_1)^3 d_{G_2}(u_2) \\ &\quad - 3{n_1}^2 d_{G_1}(u_1) d_{G_2}(u_2)^3 + 3n_2 d_{G_1}(u_1)^3 d_{G_2}(u_2)^2 + 3n_1 d_{G_1}(u_1)^2 d_{G_2}(u_2)^3 \\ &\quad - 6n_1 n_2 d_{G_1}(u_1)^2 d_{G_2}(u_2)^2). \end{aligned}$$

After simple calculations, we obtain the desired result. □

2.10. **Symmetric Difference.** The symmetric difference, of two graphs G_1 and G_2 is denoted by $G_1 \oplus G_2$, so that $|V(G_1 \oplus G_2)| = |V(G_1)| \times |V(G_2)|$ and

$$E(G_1 \oplus G_2) = \{(u_1, u_2)(v_1, v_2) : u_1v_1 \in E(G_1) \text{ or } u_2v_2 \in E(G_2) \text{ but not both}\}.$$

From definition of symmetric difference it is clear that

$$d_{G_1 \oplus G_2}(v_1, v_2) = n_2 d_{G_1}(v_1) + n_1 d_{G_2}(v_2) - 2 d_{G_1}(v_1) d_{G_2}(v_2).$$

In the following theorem we obtain the F-index of the symmetric difference of two graphs.

Theorem 11. *The F-index of $G_1 \oplus G_2$ is given by*

$$\begin{aligned} F(G_1 \oplus G_2) &= {n_2}^4 F(G_1) + {n_1}^4 F(G_2) - 8F(G_1)F(G_2) + 6n_1{n_2}^2 m_2 M_1(G_1) + 6{n_1}^2 n_2 m_1 M_1(G_2) \\ &\quad +12n_2 F(G_1)M_1(G_2) + 12n_1 F(G_2)M_1(G_1) - 12{n_2}^2 m_2 F(G_1) - 12{n_1}^2 m_1 F(G_2) \\ &\quad -12n_1 n_2 M_1(G_1)M_1(G_2). \end{aligned}$$

Proof. By definition of F-index, we have

$$\begin{aligned} F(G_1 \oplus G_2) &= \sum_{(u_1,u_2)\in V(G_1\wedge G_2)} d_{G_1\oplus G_2}(u_1,u_2)^3 \\ &= \sum_{u_1\in V(G_1)} \sum_{u_2\in V(G_2)} (n_2 d_{G_1}(u_1) + n_1 d_{G_2}(u_2) - 2d_{G_1}(u_1)d_{G_2}(u_2))^3 \\ &= \sum_{u_1\in V(G_1)} \sum_{u_2\in V(G_2)} ({n_2}^3 d_{G_1}(u_1)^3 + {n_1}^3 d_{G_2}(u_2)^3 - 8d_{G_1}(u_1)^3 d_{G_2}(u_2)^3 \\ &\quad +3n_1{n_2}^2 d_{G_1}(u_1)^2 d_{G_2}(u_2) + 3{n_1}^2 n_2 d_{G_1}(u_1)d_{G_2}(u_2)^2 - 6{n_2}^2 d_{G_1}(u_1)^3 d_{G_2}(u_2) \\ &\quad -6{n_1}^2 d_{G_1}(u_1)d_{G_2}(u_2)^3 + 12n_2 d_{G_1}(u_1)^3 d_{G_2}(u_2)^2 + 12n_1 d_{G_1}(u_1)^2 d_{G_2}(u_2)^3 \\ &\quad -12n_1 n_2 d_{G_1}(u_1)^2 d_{G_2}(u_2)^2), \end{aligned}$$

from where we obtain the desired result. □

2.11. **Splice and Link of Graphs.** Let $v_1 \in V(G_1)$ and $v_2 \in V(G_2)$ be two given vertices of G_1 and G_2, respectively. A splice of G_1 and G_2 at the vertices v_1 and v_2, denoted by $(G_1 \bullet G_2)(v_1,v_2)$, was introduced by Doslić [27], and is obtained by identifying the vertices v_1 and v_2 in the union of G_1 and G_2. The vertex set of $(G_1 \bullet G_2)(v_1,v_2)$ is given by $V((G_1 \bullet G_2)(v_1,v_2)) = [V(G_1)\backslash\{v_1\}] \cup [V(G_2)\backslash\{v_2\}] \cup \{v_{12}\}$, where we denote the vertex obtained by identifying v_1 and v_2 by v_{12}. From the construction of splice graphs it is clear that

$$d_{(G_1\bullet G_2)(v_1,v_2)}(v) = \begin{cases} d_{G_i}(v), & \text{for } v \in V(G_i) \text{ and } v \neq v_i, \\ d_{G_1}(v_1) + d_{G_2}(v_2), & \text{for } v = v_{12}. \end{cases}$$

Similarly a link of G_1 and G_2 at the vertices v_1 and v_2 is denoted by $(G_1 \sim G_2)(v_1,v_2)$ and is obtained by joining the vertices v_1 and v_2 in the union of G_1 and G_2. From the construction of link graphs it is clear that

$$d_{(G_1\sim G_2)(v_1,v_2)}(v) = \begin{cases} d_{G_i}(v), & v \in V(G_i) \text{ and } v \neq v_i, i = 1,2, \\ d_{G_i}(v) + 1, & v = v_i, i = 1,2. \end{cases}$$

In the following, we find the F-index of splice and link of two graphs G_1 and G_2 at the vertices v_1 and z.

Theorem 12. *The F-index of* $(G_1 \bullet G_2)(v_1, v_2)$ *is given by*

$$F((G_1 \bullet\ G_2)(v_1, v_2)) = F(G_1) + F(G_2) + 3d_{G_1}(v_1)d_{G_2}(v_2)(d_{G_1}(v_1) + d_{G_2}(v_2)).$$

Proof. From the definition of F-index we have

$$\begin{aligned} F((G_1 \bullet G_2)(v_1, v_2)) &= \sum_{v \in V(G_1), v \neq v_1} d_{G_1}(v)^3 + \sum_{v \in V(G_2), v \neq v_2} d_{G_2}(v)^3 + (d_{G_1}(v_1) + d_{G_2}(v_2))^3 \\ &= \sum_{v \in V(G_1)} d_{G_1}(v)^3 + \sum_{v \in V(G_2)} d_{G_2}(v)^3 + 3d_{G_1}(v_1)^2 d_{G_2}(v_2) + 3d_{G_1}(v_1)d_{G_2}(v_2)^2. \end{aligned}$$

From above we obtain the desired result after simple calculations. □

Theorem 13. *The F-index of* $(G_1 \sim\ G_2)(v_1, v_2)$ *is given by*

$$F((G_1 \sim\ G_2)(v_1, v_2)) = F(G_1) + F(G_2) + 3(d_{G_1}(v_1) + d_{G_2}(v_2)) + 3(d_{G_1}(v_1)^2 + d_{G_2}(v_2)^2) + 2.$$

Proof. From the definition of F-index, we have

$$\begin{aligned} F((G_1 \sim G_2)(v_1, v_2)) &= \sum_{v \in V(G_1), v \neq v_1} d_{G_1}(v)^3 + \sum_{v \in V(G_2), v \neq v_2} d_{G_2}(v)^3 + (d_{G_1}(v_1) + 1)^3 + (d_{G_2}(v_2) + 1)^3 \\ &= F(G_1) + F(G_2) + 3d_{G_1}(v_1) + 3d_{G_1}(v_1)^2 + 1 + 3d_{G_2}(v_2) + 3d_{G_2}(v_2)^2 + 1, \end{aligned}$$

from where the desired result follows. □

3. Conclusion

In this paper, we derive some explicit expression of the F-index of different graph operations such as union, join, Cartesian product, composition, tensor product, strong product, corona product, generalized hierarchical product, disjunction, symmetric difference, splice and link of graphs. Also we apply our results to compute the F-index for some important classes of molecular graphs and nano-structures. For further study, F-index of some other graph operations and for different composite graphs can be computed.

REFERENCES

[1] I. Gutman, N. Trinajstić, Graph theory and molecular orbitals. Total π-electron energy of alternant hydrocarbons, *Chem. Phys. Lett.*, **17**(1972), 535–538.

[2] B. Furtula, I. Gutman, A forgotten topological index, *J. Math. Chem.*, **53(4)**(2015), 1184–1190.

[3] M.H. Khalifeh, H. Yousefi-Azari, A.R. Ashrafi, The first and second Zagreb indices of some graph operations, *Discrete Appl. Math.*, **157(4)**(2009), 804–811.

[4] K.C. Das, A. Yurttas, M. Togan, A.S. Cevik and I.N. Cangul,The multiplicative Zagreb indices of graph operations, *J. Inequal. Appl.*, **2013:90**(2013), doi: 10.1186/1029-242X-2013-90.

[5] M. Azari, Sharp lower bounds on the Narumi-Katayama index of graph operations, *Appl. Math. Comput.*, **239**(2014), 409–421

[6] M. Azari, A. Iranmanesh, Computing the eccentric-distance sum for graph operations, *Discrete Appl. Math.*, **161(18)**(2013), 2827–2840.

[7] M. Azari and A. Iranmanesh, Some inequalities for the multiplicative sum Zagreb index of graph operations, *J. Math. Inequal.* **9(3)** (2015), 727–738.

[8] M. Azari and A. Iranmanesh, Chemical graphs constructed from rooted product and their Zagreb indices, *MATCH Commun. Math. Comput. Chem.* **70** (2013), 901–919.

[9] N. De, A. Pal, and S.M.A. Nayeem, The irregularity of some composite graphs, *Int. J. Appl. Comput. Math*, doi: 10.1007/s40819-015-0069-z.

[10] N. De, A. Pal, and S.M.A. Nayeem, On some bounds and exact formulae for connective eccentric indices of graphs under some graph operations *Intern. J. Combinatorics*, **2014**(2014), doi: 10.1155/2014/579257.

[11] A.R. Ashrafi, T. Doslic and A. Hamzeh, The Zagreb coindices of graph operations, *Discrete Appl. Math.*, **158**(2010), 1571–1578.

[12] M. H. Khalifeh, H. Yousefi-Azari, A. R. Ashrafi, The hyper-Wiener index of graph operations, *Comput. Math. Appl.*, **56**(2008), 1402–1407.

[13] M. Tavakoli, F. Rahbarnia and A.R. Ashrafi, Some new results on irregularity of graphs,*J. Appl. Math. Informatics*, **32**(2014), 675–685.

[14] M. Veylaki, M.J. Nikmehr and H.A. Tavallaee, The third and hyper-Zagreb coindices of some graph operations, *J. Appl. Math. Comput.*, (2015), doi: 10.1007/s12190-015-0872-z.

[15] B. Eskender and E. Vumar, Eccentric connectivity index and eccentric distance sum of some graph operations, *Trans. Comb.*, **2(1)**(2013), 103–111.

[16] L. Barrière, C. Dalfó, M. A. Fiol and M. Mitjana, The generalized hierarchical product of graphs, *Discrete Math.*, **309**(2009), 3871–3881.

[17] M. Eliasi, A. Iranmanesh, Hosoya polynomial of hierarchical product of graphs, *MATCH Commun. Math. Comput. Chem.*, **69(1)**(2013), 111–119.

[18] M. Arezoomand, B. Taeri, Zagreb indices of the generalized hierarchical product of graphs, *MATCH Commun. Math. Comput. Chem.*, **69(1)**(2013), 131–140.

[19] Z. Luo and J. Wu, Zagreb eccentricity indices of the generalized hierarchical product graphs and their applications, *J. Appl. Math.*, **2014**(2014), doi: 10.1155/2014/241712.

[20] N. De, S. M. A. Nayeem and A. Pal, Total eccentricity index of the generalized hierarchical product of graphs, *Intern. J. Appl. Comput. Math.*, (2015), doi:10.1007/s40819-014-0016-4.

[21] K. Pattabiraman and P. Kandan, Weighted PI index of corona product of graphs, *Discrete Math. Alg. Appl.*, **06**(2014), doi: 10.1142/S1793830914500554.

[22] Y. Alizadeh, A. Iranmanesh, T. Doslic and M. Azari, The edge Wiener index of suspensions, bottlenecks, and thorny graphs, *Glas. Mat. Ser. III*, **49**(2014), 1–12.

[23] Z. Yarahmadi and A. R. Ashrafi, The Szeged, Vertex PI, first and second Zagreb indices of corona product of graphs, *Filomat*, **26(3)**(2012), 467–472.

[24] N. De, On eccentric connectivity index and polynomial of thorn graph, *Appl. Math.*, **3**(2012), 931–934.

[25] N. De, Augmented eccentric connectivity index of some thorn graphs, *Intern. J. Appl. Math. Res.*, **1(4)**(2012), 671–680.

[26] N. De, A. Pal and S.M.A. Nayeem, Modified eccentric connectivity of generalized thorn graphs, *Intern. J. Computational Math.*, **2015**(2015), doi: 10.1155/2014/436140.

[27] T. Doslić, Splices, links and their degreeweighted Wiener polynomials, *Graph Theory Notes N. Y.*, **48**(2005), 47-55.

DEPARTMENT OF BASIC SCIENCES AND HUMANITIES (MATHEMATICS), CALCUTTA INSTITUTE OF ENGINEERING AND MANAGEMENT, KOLKATA, INDIA.

E-mail address: de.nilanjan@rediffmail.com

DEPARTMENT OF MATHEMATICS, ALIAH UNIVERSITY, IIA/27, NEW TOWN, KOLKATA - 700 156, INDIA.

E-mail address: nayeem.math@aliah.ac.in

DEPARTMENT OF MATHEMATICS, NATIONAL INSTITUTE OF TECHNOLOGY, DURGAPUR, INDIA.

E-mail address: anita.buie@gmail.com